

On a level-set methods for ill-posed problems with piecewise non-constant coefficients

A. De Cezaro[†]

July 12, 2012

Abstract

We investigate level-set type methods for solving ill-posed problems, with the assumption that the solutions are piecewise, but not necessarily constant functions with unknown level sets and unknown level values.

In order to get a stable approximate solutions of the inverse problem a Tikhonov-type regularization approach coupled with a level set framework is proposed. We prove existence of generalized minimizers for the Tikhonov functional. Moreover, we prove convergence and stability for regularized solutions with respect to the noise level, characterizing the level-set approach as a regularization method for inverse problems.

We also showing the applicability of the proposed level set method in some interesting inverse problems arising in elliptic PDE models.

Keywords: Level Set Methods, Regularization, Ill-Posed Problems, Piecewise Non-Constant Coefficients.

1 Introduction

Since the seminal paper of Santosa [30], level set techniques have been successful developed and has recently become a standard technique for solving inverse problems with interfaces. As references, see [35, 17, 36, 29, 27, 8, 11, 18, 5] just for cite a few.

In many applications, interfaces represent interesting physical parameters (inhomogeneities, heat conductivity between materials with different heat capacity, interface diffusion problems) across which one or more of these physical parameters change value in a discontinuous manner. The interfaces divide the domain $\Omega \subset \mathbb{R}^n$ in subdomains Ω_j , with $j = 1, \dots, k$, of different region with specific internal parameter profiles. Due to the different physical structure of each of these regions, in general, quite different mathematical models might be most appropriate for describing them. Solutions of such model represents a free boundary problem, i.e., one in which interfaces are also unknown and must be determined in addition to the solution of the governing partial differential equation [9, 11, 8, 5, 10, 26, 41, 34], in general, by a set of data obtained by indirect measurements. Applications includes image segmentation problems [10, 26, 34, 41], optimal shape designer problems [32, 5], Stefan's type problems [5], inverse potential problems [15, 13, 14], inverse conductivity/resistivity problems [24, 17, 36, 11, 9, 5] among others [8, 11, 5, 32, 18].

There is often a large variety of priors information available for determining the unknown physical parameter, whose character depends on the given application. In this article, we are interesting in inverse problems that consists in the identification of an unknown quantity $u \in D(F) \subset X$ that represents all parameter profiles inside the individual subregions of Ω , from data $y \in Y$, where X and Y are Banach spaces and $D(F)$ will be adequately specified in Section 3. In this particular case, only the interfaces

[†]Institute of Mathematics Statistics and Physics, Federal University of Rio Grande, Av. Italia km 8, 96201-900 Rio Grande, Brazil (adrianocezaro@furg.br).

between the different regions and, possibly, the unknown parameter values need to be reconstructed from the gathered data.

This process can be formally described by the operator equation

$$F(u) = y, \quad (1)$$

where $F : D(F) \subset X \rightarrow Y$ is the forward operator. In practical applications, data are obtained only by indirect measurements of the parameter. Hence, in general, exact data $y \in Y$ are not known and we have only access to noise data $y^\delta \in Y$, whose level of noise $\delta > 0$ are assumed be known *a priori* and satisfies

$$\|y^\delta - y\|_Y \leq \delta. \quad (2)$$

In particular, problems as described above are ill-posed and the accuracy of an approximated solution call for a regularization method [19, 5]. In this article we propose a Tikhonov-type regularization method coupled with a level-set approach to obtain a stable approximation of the unknown level sets and values of the piecewise solution of (1).

Many approaches, in particular level set type approaches, have been suggested for such problems previously. In [6, 7, 9, 22, 25, 30], level set approaches for identification of the unknown parameter u with distinct, but known, piecewise constant values was investigated. In [7, 10, 15], level set approaches was derived for solve inverse problems, assuming that u is defined by several distinct constant values. In both cases, one needs only to identify the level sets of u , i.e. the inverse problem reduces to a shape identification problem. On the other hand, when the level values of u are also unknown, the inverse problem becomes harder, since, we have to identify both, the level sets as well as the level values of the unknown parameter u . In this situation, the dimension of the parameter space increases by the number of unknown level values. Level set approaches to ill- posed problems with unknown constant level values appear before in [14, 13, 32, 33, 34]. For level set regularization properties of the approximated solution for inverse problems see [15, 13, 14, 22, 4]. However, regularization theory for inverse problems where the components of the parameter u are variable and has jumps of discontinuities was not well investigate. Some early results in the numerical implementation of such inverse problems was consider before in [11].

In this article, we propose a level set type regularization method to ill-posed problems whose solution is composed by piecewise components not necessarily constants. These type of problems, where the parameters are variable and discontinuous, arise in many problems as heat conduction between materials of different heat capacity and conductivity, interface diffusion processes and many other types of physical problems where modeling components are related with embedded boundaries. See, for example [8, 11, 18, 3] and references.

The advantage of our proposed level set method *vis-a-vis* the approaches before is that, we prove regularization properties for the approximated solution of the inverse problem model by (1) where the parameter is a non-constant piecewise solutions. Therefore such level set type method is different of those in the literature since we are able to apply this method to problems whose solutions depends of moving interfaces and non-constant parameters. In terms of applications, the presented regularization approach can handle with more kinds of flexibility and interesting physical problems, where de components of the interesting parameter is not necessarily homogeneous [18, 14, 13, 32, 33, 34, 3]. As a benchmark problem we analyze two inverse problem model by elliptic PDE's with discontinuous and variable interesting coefficient.

The paper is outlined as follows: In Section 2, we formulate the Tikhonov functional based on the level-set framework. In Section 3, we present the general assumptions needed in this article and the definition of the set of admissible solutions. In Subsection 3.1, we prove relevant properties about the admissible set of solutions, in particular, convergence in suitable topologies. Relevant properties of the penalization functional is given in Subsection 3.2. In Section 4, we prove that the proposed method is a regularization method to inverse problem, i.e., we prove that the minimizers of the proposed Tikhonov functional are stable and convergent with respect to the noise level in the data. In Section 5, a smooth functional is proposed to approximate minimizers of the Tikhonov functional defined in the admissible

set of solutions. We provide approximation properties and optimality condition for the minimizers of the smooth Tikhonov functional. In Section 6, we present an application of the proposed framework to solve some interesting inverse elliptic problem with variable coefficient. Conclusions and future directions are presented in Section 7.

2 The Level-set Formulation

Our starting point is the assumption that the parameter u in (1) assume two unknown functional values, i.e., $u(x) \in \{\psi^1(x), \psi^2(x)\}$ a.e. in $\Omega \subset \mathbb{R}^d$, where Ω is a bounded set. More specifically, we assume the existence of a measurable set $D \subset \subset \Omega$, with $0 < |D| < |\Omega|$, such that $u(x) = \psi^1(x)$ if $x \in D$ and $u(x) = \psi^2(x)$ if $x \in \Omega/D$.

With this framework, the inverse problem that we are interesting in this article is the stable identification of both, the shape of D and the value function $\psi^j(x)$ for x belongs to D and to Ω/D , respectively, from observation of the data $y^\delta \in Y$.

We remark that, if $\psi^1(x) = c^1$ and $\psi^2(x) = c^2$ with c^1 and c^2 unknown constants values, the problem of identify u was rigorously studies before in [14]. Moreover, many other approaches to this case appear in the literature. See [14, 7, 6, 5] and references. Recently, in [13], a L^2 level set approach to identify the level and constant contrast was investigated.

Our approach differs from the level set methods proposed in [13, 14], by consider also the identification of variable unknown level of the parameter u . In this situation, many topological difficulties appears in order to have a tractable definition of an admissible set of parameters (see Definition 1 below). Generalization to problems with more than tow levels following by the techniques derived in [15], applied to this approach. As observed before, the present level set approach is a rigorously derivation of a regularization strategies for identification of the shape and non-constant levels of discontinuous parameters. Therefore, it can be applied to physical problems modeled be embedded boundaries that whose components are not necessarily piecewise constant [22, 5, 15, 13, 14].

In many interesting applications, the equation (1) is ill-posed. Therefore, in order to obtain a stable approximate solution, some regularization method will be applied. We propose a regularization method by: First, introduce a parametrization on the parameter space, using a level set function ϕ that belongs to $H^1(\Omega)$. Note that, we can identify the distinct level sets of the function $\phi \in H^1(\Omega)$ with the definition of the Heaviside projector

$$H : H^1(\Omega) \longrightarrow L^\infty(\Omega)$$

$$\phi \longmapsto H(\phi) := \begin{cases} 1 & \text{if } \phi(x) > 0, \\ 0 & \text{other else.} \end{cases}$$

Now, from the framework introduced above, a solution of (1) can be represented as

$$u(x) = \psi^1(x)H(\phi) + \psi^2(x)(1 - H(\phi)) =: P(\phi, \psi^1, \psi^2)(x). \quad (3)$$

With this notation, we are able to determine the shapes of D as $\{x \in \Omega; \phi(x) > 0\}$ and Ω/D as $\{x \in \Omega; \phi(x) < 0\}$.

The functional level values $\psi^1(x), \psi^2(x)$ are also assumed be unknown and they needed to be determined as well.

Assumption 1. We assume that $\psi^1, \psi^2 \in \mathbb{B} := \{f : f \text{ is measurable and } f(x) \in [m, M], \text{ a.e. in } \Omega\}$, for some constant values m, M .

Remark 1. We remark that, $f \in \mathbb{B}$ implies that $f \in L^\infty(\Omega)$. Since Ω is bounded $f \in L^1(\Omega)$. Moreover,

$$\int_{\Omega} f(x) \nabla \cdot \varphi(x) dx \leq |M| \int_{\Omega} |\nabla \cdot (\varphi)(x)| dx \leq |M| \|\nabla \cdot \varphi\|_{L^1(\Omega)}, \quad \forall \varphi \in C_0^1(\Omega, \mathbb{R}^n).$$

Hence $f \in \text{BV}(\Omega)$.

Note that, in the case that ψ^1 and ψ^2 assumes tow distinct constant values (as covered by the analysis done in [5, 14, 13] and references) the assumptions above are satisfied. Hence, the level set approach proposed here generalize the regularization theory developed in [14, 13].

From (3), the inverse problem in (1), with data given as in (2), can be abstractly written as the operator equation

$$F(P(\phi, \psi^1, \psi^2)) = y^\delta. \quad (4)$$

Since, an approximate solution (ϕ, ψ^1, ψ^2) of (4) is obtained, a corresponding solution of (1) can be computed using equation (3).

Therefore, for obtain a regularized approximate solutions to (4), we shell consider the least square approach combined with a regularization term i.e., minimizing the Tikhonov functional

$$\hat{\mathcal{G}}_\alpha(\phi, \psi^1, \psi^2) := \|F(P(\phi, \psi^1, \psi^2)) - y^\delta\|_Y^2 + \alpha \left\{ \beta_1 |H(\phi)|_{\text{BV}} + \beta_2 \|\phi - \phi_0\|_{H^1(\Omega)}^2 + \beta_3 \sum_{j=1}^2 |\psi^j - \psi_0^j|_{\text{BV}} \right\}, \quad (5)$$

where, ϕ_0 and ψ_0^j represents some *a priori* information about the true solution u^* of (1). The parameter $\alpha > 0$ play the rule of regularization parameter and the values of $\beta_i, i = 1, 2, 3$ act as scaling factors. In other words, $\beta_i, i = 1, 2, 3$ need to be chosen *a priori*, but independent of the noise level δ . In practical, $\beta_i, i = 1, 2, 3$ can be chosen in order to represent *a priori* knowledge of features the of parameter solution u and/or to improve the numerical algorithm. A complete discussion about how to choose $\beta_i, i = 1, 2, 3$ are provided in [13, 14, 15].

The regularization strategy in this context is based on $TV - H^1 - TV$ penalization. The term on H^1 -norm act simultaneous as a control on the size of the norm of the level set function and a regularization on the space H^1 . The term on BV is a variational measure of $H(\phi)$. It is well known for penalizing the length of the Hausdorff measure of the boundary of the set $\{x : \phi(x) > 0\}$ (see [21, Chapter 5] for details). Finally, the last term on BV is a variational measure of ψ^j that acts as a regularization term on the set $\mathbb{B}$. This Tikhonov functional extends the ones proposed in [14, 15, 6, 7, 32] (based on $TV - H^1$ penalization).

Existence of minimizers for the functional (5), in the $H^1 \times \mathbb{B}^2$ topology do not follows by direct arguments, since, the operator P is not necessary continuous in this topology. Indeed, is $\psi^1 = \psi^2 = \psi$ a continuous function at the contact region, than $P(\phi^1, \psi^2, \psi) = \psi$ is continuous and the standard Tikhonov regularization theory to inverse problem holds true [19]. On the other hand, in the interesting case where ψ^1 and ψ^2 represents the level of the jumps in the parameter u , the analysis became more complicated and we need a definition of generalized minimizers (see Definition 1) in order to handle with the difficulties.

The advantage of our approach in relation to [5, 14, 22, 15] is that we are able to obtain regularize solutions to inverse problems with piecewise solutions that are more general that those covered by the regularization methods proposed before. In particular, we propose a level set regularization method to regularize piecewise non-constant solutions of ill-posed problems. Therefore, the approach studied here can be applied to a variate of physical problems of interesting as heat conduction between materials of different heat capacity and conductivity and interface diffusion processes, where the coefficient are variable and discontinuous [8, 11, 3].

We remark that, level set type methods was used for obtain solutions of elliptic problems with discontinuous and variable coefficients [11]. However, level set regularization theory for solve inverse problems [13, 15, 14] for recover the shape and the values of variable discontinuous coefficient are unknown be the author.

The principal difference between our approach *vis a vis* the other ones before [14, 33, 32, 34, 30, 25, 22, 9, 6, 7], is that we introduce a level set type regularization for recover the shape and the values of variable discontinuous coefficient. In this framework, the topologies needed to guaranties the existence of a minimizer (in a generalized sense) of the Tikhonov functional (5) is quite complicate and differ in some key points of [13, 14]. In these particular approach, the definition of generalized minimizer are quite different of others before [14, 22, 15]. As a consequence, the arguments used to prove the well-posed of the Tikhonov functional (5), the stability and convergence of the regularized solutions of the inverse problem (1) are quite complicated and needed significant improvements (see Definition 1).

3 Generalized Minimizers

As already observed in [22], if $D \subset \Omega$ with $\mathcal{H}^{n-1}(\partial D) < \infty$ where $\mathcal{H}^{n-1}(S)$ denotes the (n-1)-dimensional Hausdorff-measure of the set S , then the Heaviside operator H maps $H^1(\Omega)$ into the set

$$\mathcal{V} := \{\chi_D; D \subset \Omega \text{ measurable, } : \mathcal{H}^{n-1}(\partial D) < \infty\}.$$

Therefore, the operator P in (3) maps $H^1(\Omega) \times \mathbb{B}^2$ into the admissible parameter set

$$D(F) := \{u = q(v, \psi^1, \psi^2); v \in \mathcal{V} \text{ and } \psi^1, \psi^2 \in \mathbb{B}\},$$

where

$$q : \mathcal{V} \times \mathbb{B}^2 \ni (v, \psi^1, \psi^2) \mapsto \psi^1 v + \psi^2 (1 - v) \in BV(\Omega).$$

Consider the model problem described as in the introduction. In this article, we assume that:

(A1) $\Omega \subseteq \mathbb{R}^n$ is bounded with piecewise C^1 boundary $\partial\Omega$.

(A2) The operator $F : D(F) \subset L^1(\Omega) \rightarrow Y$ is continuous on $D(F)$ with respect to the $L^1(\Omega)$ -topology.

(A3) ε, α and $\beta_j, j = 1, 2, 3$ denote positive parameters.

(A4) Equation (1) has a solution, i.e. there exists $u_* \in D(F)$ satisfying $F(u_*) = y$ and a function $\phi_* \in H^1(\Omega)$ satisfying $|\nabla \phi_*| \neq 0$, in a neighborhood of $\{\phi_* = 0\}$ such that $H(\phi_*) = z_*$, for some $z_* \in \mathcal{V}$. Moreover, there exist functional values $\psi_*^1, \psi_*^2 \in \mathbb{B}$ such that $q(z_*, \psi_*^1, \psi_*^2) = u_*$.

For each $\varepsilon > 0$, we define an smooth approximation to the operator P by

$$P_\varepsilon(\phi, \psi^1, \psi^2) := \psi^1 H_\varepsilon(\phi) + \psi^2 (1 - H_\varepsilon(\phi)), \quad (6)$$

where H_ε is the smooth approximation to H describe by

$$H_\varepsilon(t) := \begin{cases} 1 + t/\varepsilon & \text{for } t \in [-\varepsilon, 0] \\ H(t) & \text{for } t \in \mathbb{R} \setminus [-\varepsilon, 0] \end{cases}.$$

Remark 2. It is worth noting that, for any $\phi_k \in H^1(\Omega)$, $H_\varepsilon(\phi_k)$ belongs to $L^\infty(\Omega)$ and satisfies $0 \leq H_\varepsilon(\phi_k) \leq 1$ a.e. in Ω , for all $\varepsilon > 0$. Moreover, taking in account that $\psi^j \in \mathbb{B}$, follows that the operators q and P_ε , as above, are well defined.

In order to guarantee existence of a minimizer of $\mathcal{G}_\alpha$ defined in (5) in the space $H^1(\Omega) \times \mathbb{B}^2$, we need introduce a suitable topology such that the functional $\mathcal{G}_\alpha$ has closed graphic. Therefore, the concept of generalized minimizers (compare with [15, 22]) in this paper is:

Definition 1. Let the operators H, P, H_ε and P_ε be defined as above and positive parameters α, β_j and ε given in the general assumption.

A **quadruple** $(z, \phi, \psi^1, \psi^2) \in L^\infty(\Omega) \times H^1(\Omega) \times BV(\Omega)^2$ is called **admissible** when:

- a) There exists a sequence $\{\phi_k\}$ of $H^1(\Omega)$ -functions satisfying $\lim_{k \rightarrow \infty} \|\phi_k - \phi\|_{L^2(\Omega)} = 0$.
- b) There exists a sequence $\{\varepsilon_k\} \in \mathbb{R}^+$ converging to zero such that $\lim_{k \rightarrow \infty} \|H_{\varepsilon_k}(\phi_k) - z\|_{L^1(\Omega)} = 0$.
- c) There exist sequences $\{\psi_k^1\}_{k \in \mathbb{N}}$ and $\{\psi_k^2\}_{k \in \mathbb{N}}$ belongs to $BV \cap C^\infty(\Omega)$ such that

$$|\psi_k^j|_{BV} \longrightarrow |\psi^j|_{BV}, \quad j = 1, 2.$$

- d) A **generalized minimizer** of $\hat{\mathcal{G}}_\alpha$ is considered to be any admissible quadruple $(z, \phi, \psi^1, \psi^2)$ minimizing

$$\mathcal{G}_\alpha(z, \phi, \psi^1, \psi^2) := \left\| F(q(z, \psi^1, \psi^2)) - y^\delta \right\|_Y^2 + \alpha R(z, \phi, \psi^1, \psi^2) \quad (7)$$

on the set of admissible quadruples. Here the functional R is defined by

$$R(z, \phi, \psi^1, \psi^2) = \rho(z, \phi) + \beta_3 \sum_{j=1}^2 |\psi^j - \psi_0^j|_{\text{BV}} \quad (8)$$

and the functional ρ is defined as

$$\rho(z, \phi) := \inf \left\{ \liminf_{k \rightarrow \infty} \left[\beta_1 |H_{\varepsilon_k}(\phi_k)|_{\text{BV}} + \beta_2 \|\phi_k - \phi_0\|_{H^1(\Omega)}^2 \right] \right\}. \quad (9)$$

The infimum in (9) is taken over all sequences $\{\varepsilon_k\}$ and $\{\phi_k\}$ characterizing $(z, \phi, \psi^1, \psi^2)$ as an admissible quadruple.

The convergence $|\psi_k^j|_{\text{BV}} \rightarrow |\psi^j|_{\text{BV}}$ in Item c) in Definition 1 is in the sense of variation measure [21, Chapter 5]. The incorporation of item c) in the Definition 1 implies in the existence of the Γ -limit of sequences of admissible quadruples [22, 1]. This appears in the proof of Lemmas 4, 5 and 8, where we proof that the set of admissible quadruple are closed in the defined topology (see Lemmas 4 and 5) and the weak lower semi-continuity of the regularization functional R (see Lemma 8). The identification of non-constant level values ψ^j implies in a different definition of admissible quadruple. As a consequence the arguments in the proof of regularization properties of the level set approach and the applications are the principal and novel difference between our definition of admissible quadruple in the present article and one that appears in [16, 13, 14, 22].

Remark 3. For $j = 1, 2$ let $\psi^j \in \mathbb{B} \cap C^\infty(\Omega)$, $\phi \in H^1(\Omega)$ be such that $|\nabla \phi| \neq 0$ in a neighborhood of the level set $\{\phi(x) = 0\}$ and $H(\phi) = z \in \mathcal{V}$. For each $k \in \mathbb{N}$ set $\psi_k^j = \psi^j$ and $\phi_k = \phi$. Then, for all sequences of $\{\varepsilon_k\}_{k \in \mathbb{N}}$ of positive numbers converging to zero, we have

$$\begin{aligned} \|H_{\varepsilon_k}(\phi_k) - z\|_{L^1(\Omega)} &= \|H_{\varepsilon_k}(\phi_k) - H(\phi)\|_{L^1(\Omega)} = \int_{(\phi)^{-1}[-\varepsilon_k, 0]} \left| 1 - \frac{\phi}{\varepsilon_k} \right| dx \\ &\leq \int_{-\varepsilon_k}^0 \int_{(\phi)^{-1}(\tau)} 1 d\tau \leq \text{meas}\{(\phi)^{-1}(\tau)\} \int_{-\varepsilon_k}^0 1 dt \rightarrow 0. \end{aligned}$$

Here, we use the fact that $|\nabla \phi| \neq 0$ in a neighborhood of $\{\phi = 0\}$ implies that ϕ is a local diffeomorphism together with a co-area formula [21, Chapter 4]. Moreover, $\{\psi_k^j\}_{k \in \mathbb{N}}$ in $\mathbb{B} \cap C^\infty(\Omega)$ satisfying Definition 1 item c).

Hence, $(z, \phi, \psi^1, \psi^2)$ is an admissible quadruple. In particular, we conclude from the general assumption above that the set of admissible quadruple satisfying $F(u) = y$ is not empty.

3.1 Relevant Properties of Admissible Quadruples

Our first result will be to prove continuities properties of operators P_ε , H_ε and q in suitable topologies. Such results will be necessary in the subsequent analysis.

We start with a auxiliary lemma that is well known (see for example [16]). We present it here for sake of completeness.

Lemma 2. Let Ω be a measurable subset of $\mathbb{R}^n$ with finite measure.

If $(f_k) \in \mathbb{B}$ is a convergent sequence in $L^p(\Omega)$ for some p , $1 \leq p < \infty$, then it is a convergent sequence in $L^p(\Omega)$ for all $1 \leq p < \infty$.

In particular Lemma 2 holds for the sequence $z_k := H_\varepsilon(\phi_k)$.

Proof. See [16, Lemma 2.1]. □

Next two lemmas are auxiliary results in order to understand the definition of the set of admissible quadruple.

Lemma 3. Let Ω as in the assumption (A1) and $j = 1, 2$.

- (i) Let $\{z_k\}_{k \in \mathbb{N}}$ be a sequence in $L^\infty(\Omega)$ with $z_k \in [m, M]$ a.e. converging in the $L^1(\Omega)$ -norm to some element z and $\{\psi_k^j\}_{k \in \mathbb{N}}$ be a sequence in $\mathbb{B}$ converging in the BV-norm to some $\psi^j \in \mathbb{B}$. Then $q(z_k, \psi_k^1, \psi_k^2)$ converges to $q(z, \psi^1, \psi^2)$ in $L^1(\Omega)$.
- (ii) Let $(z, \phi) \in L^1(\Omega) \times H^1(\Omega)$, be such that $H_\varepsilon(\phi) \rightarrow z$ in $L^1(\Omega)$ as $\varepsilon \rightarrow 0$ and let $\psi^1, \psi^2 \in \mathbb{B}$. Then $P_\varepsilon(\phi, \psi^1, \psi^2) \rightarrow q(z, \psi^1, \psi^2)$ in $L^1(\Omega)$ as $\varepsilon \rightarrow 0$.
- (iii) Given $\varepsilon > 0$, let $\{\phi_k\}_{k \in \mathbb{N}}$ be a sequence in $H^1(\Omega)$ converging to $\phi \in H^1(\Omega)$ in the L^2 -norm. Then $H_\varepsilon(\phi_k) \rightarrow H_\varepsilon(\phi)$ in $L^1(\Omega)$, as $k \rightarrow \infty$. Moreover, if $\{\psi_k^j\}_{k \in \mathbb{N}}$ are sequences in $\mathbb{B}$, converging to some ψ^j in $\mathbb{B}$, with respect to the $L^1(\Omega)$ -norm, then $q(H_\varepsilon(\phi_k), \psi_k^1, \psi_k^2) \rightarrow q(H_\varepsilon(\phi), \psi^1, \psi^2)$ in $L^1(\Omega)$, as $k \rightarrow \infty$.

Proof. Since Ω is assumed to be bounded, we have $L^\infty(\Omega) \subset L^1(\Omega)$ and $BV(\Omega)$ is continuous embedding in $L^2(\Omega)$ [21]. To prove (i), notice that

$$\begin{aligned} \|q(z_k, \psi_k^1, \psi_k^2) - q(z, \psi^1, \psi^2)\|_{L^1(\Omega)} &= \|\psi_k^1 z_k + \psi_k^2(1 - z_k) - \psi^1 z - \psi^2(1 - z)\|_{L^1(\Omega)} \\ &\leq \|z_k\|_{L^\infty(\Omega)} \|\psi_k^1 - \psi^1\|_{L^1(\Omega)} + \|\psi^1\|_{L^2(\Omega)} \|z_k - z\|_{L^2(\Omega)} \\ &\quad + \|1 - z_k\|_{L^\infty(\Omega)} \|\psi_k^2 - \psi^2\|_{L^1(\Omega)} + \|\psi^2\|_{L^2(\Omega)} \|z_k - z\|_{L^2(\Omega)} \xrightarrow{k \rightarrow \infty} 0. \end{aligned}$$

Here we use Lemma 2 in order to guarantee the convergence of z_k to z in $L^2(\Omega)$.

Assertion (ii) follows with similar arguments and the fact that $H_\varepsilon(\phi) \in L^\infty(\Omega)$ for all $\varepsilon > 0$.

As $\|H_\varepsilon(\phi_k) - H_\varepsilon(\phi)\|_{L^1(\Omega)} \leq \varepsilon^{-1} \sqrt{\text{meas}(\Omega)} \|\phi_k - \phi\|_{L^2(\Omega)}$ the first part of assertion (iii) follows. The second part of the assertion (iii) holds by a combination of the inequality above and steps in the prove of assertion (i). $\square$

Lemma 4. Let $\{\psi_k^j\}_{k \in \mathbb{N}}$ be a sequence of functions satisfying the Definition 1 converging in $L^1(\Omega)$ to some ψ^j , for $j = 1, 2$. Then ψ^j also satisfies the Definition 1.

Sketch of the proof.

Let $k \in \mathbb{N}$ and $j = 1, 2$. Since ψ_k^j satisfies Definition 1, $\psi_k^j \in \text{BV}$. From [21, Theorem 2, pg 172] there exist sequences $\{\psi_{k,l}^j\}_{l \in \mathbb{N}}$ in $\text{BV} \times C^\infty(\Omega)$ such that

$$\psi_{k,l}^j \xrightarrow{l \rightarrow \infty} \psi_k^j \text{ in } L^1(\Omega) \text{ and } |\psi_{k,l}^j|_{\text{BV}} \xrightarrow{l \rightarrow \infty} |\psi_k^j|_{\text{BV}}.$$

In particular, for the subsequence $\{\psi_{k,l(k)}^j\}_{k \in \mathbb{N}}$ follow that

$$\psi_{k,l(k)}^j \xrightarrow{k \rightarrow \infty} \psi^j \text{ in } L^1(\Omega) \text{ and } |\psi_{k,l(k)}^j|_{\text{BV}} \xrightarrow{k \rightarrow \infty} |\psi^j|_{\text{BV}}. \quad (10)$$

Moreover, by assumption $\psi^j \in L^1(\Omega)$. From the lower semi-continuity of variational measure (see [21, Theorem 1 pg. 172]), equation (10) and definition of BV space follows that $\psi^j \in \text{BV}$. $\square$

In the next lemma we prove that the set of admissible quadruple is closed with respect the $L^1(\Omega) \times L^2(\Omega) \times (L^1(\Omega))^2$ topology.

Lemma 5. Let $(z_k, \phi_k, \psi_k^1, \psi_k^2)$ be a sequence of admissible quadruples converging in $L^1(\Omega) \times L^2(\Omega) \times (L^1(\Omega))^2$ to some $(z, \phi, \psi^1, \psi^2)$, with $\phi \in H^1(\Omega)$. Then, $(z, \phi, \psi^1, \psi^2)$ is also an admissible quadruple.

Sketch of the proof. Let $k \in \mathbb{N}$. Since $(z_k^1, \phi_k^1, \psi_k^1, \psi_k^2)$ is an admissible quadruple, it follows from Definition 1 that there exist sequences $\{\phi_{k,l}\}_{l \in \mathbb{N}}$ in $H^1(\Omega)$, $\{\psi_{k,l}^1\}_{l \in \mathbb{N}}$, $\{\psi_{k,l}^2\}_{l \in \mathbb{N}}$ in $\text{BV} \times C^\infty(\Omega)$ and a correspondent sequence $\{\varepsilon_k^l\}_{l \in \mathbb{N}}$ converging to zero such that

$$\phi_{k,l} \xrightarrow{l \rightarrow \infty} \phi_k \text{ in } L^2(\Omega), \quad H_{\varepsilon_k^l}(\phi_{k,l}) \xrightarrow{l \rightarrow \infty} z_k \text{ in } L^1(\Omega) \quad \text{and} \quad |\psi_{k,l}^j|_{\text{BV}} \xrightarrow{l \rightarrow \infty} |\psi_k^j|_{\text{BV}}, j = 1, 2.$$

Define the monotone increasing function $\tau : \mathbb{N} \rightarrow \mathbb{N}$ such that, for every $k \in \mathbb{N}$ it holds

$$\varepsilon_k^{\tau(k)} \leq \frac{1}{2} \varepsilon_{k-1}^{\tau(k-1)}, \quad \|\phi_{k,\tau(k)} - \phi_k\|_{L^2(\Omega)} \leq \frac{1}{k}, \quad \|H_{\varepsilon_k^{\tau(k)}}(\phi_{k,\tau(k)}) - z_k\|_{L^1(\Omega)} \leq \frac{1}{k}, \quad |\psi_{k,\tau(k)}^j|_{\text{BV}} \longrightarrow |\psi_k^j|_{\text{BV}}, \quad j = 1, 2. \quad (11)$$

Hence, for each $k \in \mathbb{N}$

$$\begin{aligned} \|\phi - \phi_{k,\tau(k)}\|_{L^2(\Omega)} &\leq \|\phi - \phi_k\|_{L^2(\Omega)} + \|\phi_{k,\tau(k)} - \phi_k\|_{L^2(\Omega)} \\ \|z - H_{\varepsilon_k^{\tau(k)}}(\phi_{k,\tau(k)})\|_{L^1(\Omega)} &\leq \|z - z_k\|_{L^1(\Omega)} + \|H_{\varepsilon_k^{\tau(k)}}(\phi_{k,\tau(k)}) - z_k\|_{L^1(\Omega)}. \end{aligned}$$

From (11),

$$\lim_{k \rightarrow \infty} \|\phi - \phi_{k,\tau(k)}\|_{L^2(\Omega)} = 0, \quad \lim_{k \rightarrow \infty} \|z - H_{\varepsilon_k^{\tau(k)}}(\phi_{k,\tau(k)})\|_{L^1(\Omega)} = 0. \quad (12)$$

Moreover, with the same arguments as Lemma 4, follows that

$$|\psi_{k,\tau(k)}^j|_{\text{BV}} \rightarrow |\psi^j|_{\text{BV}}, \quad j = 1, 2,$$

and $\psi^j \in \text{BV}(\Omega)$. Therefore, it remains to prove that $(z, \phi, \psi^1, \psi^2)$ is an admissible quadruple. From Definition 1 and Lemma 4, it is enough to prove that $z \in L^\infty(\Omega)$. If this is not the case, there would exist a $\Omega' \subset \Omega$ with $|\Omega'| > 0$ and $\gamma > 0$ such that $z(x) > 1 + \gamma$ in Ω' (the other case: $z(x) < -\gamma$ is analogous). Since $(H_{\varepsilon_k^{\tau(k)}}(\phi_{k,\tau(k)}))(x) \in [0, 1]$ a.e. in Ω for $k \in \mathbb{N}$ (see remark after Definition 1), we would have

$$\|z - H_{\varepsilon_k^{\tau(k)}}(\phi_{k,\tau(k)})\|_{L^1(\Omega)} \geq \|z - H_{\varepsilon_k^{\tau(k)}}(\phi_{k,\tau(k)})\|_{L^1(\Omega')} \geq \gamma |\Omega'|, \quad k \in \mathbb{N},$$

contradicting the second limit in (12). $\square$

3.2 Relevant Properties of the Penalization Functional

In next lemmas, we verify properties of the functional R which are fundamental for the convergence analysis in Section 4. In particular, these properties implies that the level sets of $\mathcal{G}_\alpha$ are compact in the set of admissible quadruple, i.e., $\mathcal{G}_\alpha$ assume a minimizer on this set. First, we prove a lemma that simplify the functional R in (8). Here we present the sketch of the proof. For more details, see the arguments in [14, Lemma 3].

Lemma 6. *Let $(z, \phi, \psi^1, \psi^2)$ be an admissible quadruple. Then, there exists sequences $\{\varepsilon_k\}_{k \in \mathbb{N}}$, $\{\phi_k\}_{k \in \mathbb{N}}$ and $\{\psi_k^j\}_{k \in \mathbb{N}}$ as in the Definition 1, such that*

$$R(z, \phi, \psi^1, \psi^2) = \lim_{k \rightarrow \infty} \left\{ \beta_1 |H_{\varepsilon_k}(\phi_k)|_{\text{BV}} + \beta_2 \|\phi_k - \phi_0\|_{H^1(\Omega)}^2 + \beta_3 \sum_{j=1}^2 |\psi_k^j - \psi_0^j|_{\text{BV}} \right\}. \quad (13)$$

Sketch of the proof. For each $l \in \mathbb{N}$, the definition of R (see Definition 1) guaranties the existence of sequences ε_k^l , $\{\phi_{k,l}^j\} \in H^1(\Omega)$ and $\{\psi_{k,l}^j\} \in \mathbb{B}$ such that

$$R(z, \phi, \psi^1, \psi^2) = \lim_{l \rightarrow \infty} \left\{ \liminf_{k \rightarrow \infty} \left\{ \beta_1 |H_{\varepsilon_k^l}(\phi_{k,l})|_{\text{BV}} + \beta_2 \|\phi_{k,l} - \phi_0\|_{H^1(\Omega)}^2 \right\} + \beta_3 \sum_{j=1}^2 |\psi_{k,l}^j - \psi_0^j|_{\text{BV}} \right\}.$$

Now a similar extraction of subsequences as in Lemma 5 complete the proof. $\square$

In the following, we prove tow lemmas that are essential to the proof of well posedness of the Tikhonov functional (5).

Lemma 7. *The functional R in (8) is coercive on the set of admissible quadruples. In other words, given any admissible quadruple $(z, \phi, \psi^1, \psi^2)$ we have*

$$R(z, \phi, \psi^1, \psi^2) \geq \left(\beta_1 |z|_{\text{BV}} + \beta_2 \|\phi - \phi_0\|_{H^1(\Omega)}^2 + \beta_2 \sum_{j=1}^2 |\psi^j - \psi_0^j|_{\text{BV}} \right).$$

Sketch of the proof. Let $(z, \phi, \psi^1, \psi^2)$ be an admissible quadruple. From [15, Lemma 4], it follows that

$$\rho(z, \phi) \geq (\beta_1 |z|_{\text{BV}} + \beta_2 \|\phi - \phi_0\|_{H^1(\Omega)}^2). \quad (14)$$

Now, from (14) and the definition of R in (8), we have

$$(\beta_1 |z|_{\text{BV}} + \beta_2 \|\phi - \phi_0\|_{H^1(\Omega)}^2 + \beta_3 \sum_{j=1}^2 |\psi^j - \psi_0^j|_{\text{BV}}) \leq \rho(z, \phi) + \beta_3 \sum_{j=1}^2 |\psi^j - \psi_0^j|_{\text{BV}} = R(z, \phi, \psi^1, \psi^2),$$

concluding the proof. $\square$

Lemma 8. *The functional R in (8) is weak lower semi-continuous on the set of admissible quadruples, i.e. given a sequence $\{(z_k, \phi_k, \psi_k^1, \psi_k^2)\}$ of admissible quadruples such that $z_k \rightarrow z$ in $L^1(\Omega)$, $\phi_k \rightharpoonup \phi$ in $H^1(\Omega)$, $\psi_k^j \rightarrow \psi^j$ in $L^1(\Omega)$, for some admissible quadruple $(z, \phi, \psi^1, \psi^2)$, then*

$$R(z, \phi, \psi^1, \psi^2) \leq \liminf_{k \in \mathbb{N}} R(z_k, \phi_k, \psi_k^1, \psi_k^2).$$

Proof. The functional $\rho(z, \phi)$ is weak lower semi-continuous cf. [15, Lemma 5]. As $\psi_k^j \in \text{BV}$ follows from [21, Theorem 2 pg 172] that there exist sequences $\{\psi_{k,l}^j\} \in \text{BV} \cap C^\infty(\Omega)$ such that $\|\psi_{k,l}^j - \psi_k^j\|_{L^1(\Omega)} \leq \frac{1}{l}$. From a diagonal argument, we can extract a subsequence $\{\psi_{k,l(k)}^j\}$ of $\{\psi_{k,l}^j\}$ such that $\{\psi_{k,l(k)}^j\} \rightarrow \psi^j$ in $L^1(\Omega)$ as $k \rightarrow \infty$. Let $\xi \in C_c^1(\Omega, \mathbb{R}^n)$, $|\xi| \leq 1$. Then, from [21, Theorem 1 pg 167], it follows that

$$\begin{aligned} \int_{\Omega} \psi^j \nabla \cdot \xi dx &= \lim_{k \rightarrow \infty} \int_{\Omega} \psi_{k,l(k)}^j \nabla \cdot \xi dx = \lim_{k \rightarrow \infty} \left[\int_{\Omega} (\psi_{k,l(k)}^j - \psi_k^j) \nabla \cdot \xi dx + \int_{\Omega} \psi_k^j \nabla \cdot \xi dx \right] \\ &\leq \lim_{k \rightarrow \infty} \left[\|\psi_{k,l(k)}^j - \psi_k^j\|_{L^1(\Omega)} \|\nabla \cdot \xi\|_{L^\infty(\Omega)} |\Omega| - \int_{\Omega} \xi \cdot \sigma_k d|\psi_k^j|_{\text{BV}} \right] \leq \liminf_{k \rightarrow \infty} |\psi_k^j|_{\text{BV}}. \end{aligned}$$

Thus, from the definition of $|\cdot|_{\text{BV}}$ (see [21]), we have

$$|\psi^j|_{\text{BV}} = \sup \left\{ \int_{\Omega} \psi^j \nabla \cdot \xi dx ; \xi \in C_c^1(\Omega, \mathbb{R}^n), |\xi| \leq 1 \right\} \leq \liminf_{k \rightarrow \infty} |\psi_k^j|_{\text{BV}}.$$

Now, the lemma follows from the fact that the functional R in (8) is a linear combination of lower semi-continuous functionals. $\square$

4 Convergence Analysis

In the following, we consider any positive parameter $\alpha, \beta_j, j = 1, 2, 3$ as in the general assumption to this article. First, we prove that the functional $\mathcal{G}_\alpha$ in (7) is well posed.

Theorem 9 (Well-Posedness). *The functional $\mathcal{G}_\alpha$ in (7) attains minimizers on the set of admissible quadruples.*

Proof. Notice that, the set of admissible quadruples is not empty, since $(0, 0, 0, 0)$ is admissible. Let $\{(z_k, \phi_k, \psi_k^1, \psi_k^2)\}$ be a minimizing sequence for $\mathcal{G}_\alpha$, i.e. a sequence of admissible quadruples satisfying $\mathcal{G}_\alpha(z_k, \phi_k, \psi_k^1, \psi_k^2) \rightarrow \inf \mathcal{G}_\alpha \leq \mathcal{G}_\alpha(0, 0, 0, 0) < \infty$. Then, $\{\mathcal{G}_\alpha(z_k, \phi_k, \psi_k^1, \psi_k^2)\}$ is a bounded sequence of real numbers. Therefore, $\{(z_k, \phi_k, \psi_k^1, \psi_k^2)\}$ is uniformly bounded in $\text{BV} \times H^1(\Omega) \times \text{BV}^2$. Thus, from the Sobolev Embedding Theorem [2, 21], we guarantee the existence of a subsequence (denoted again by $\{(z_k, \phi_k, \psi_k^1, \psi_k^2)\}$) and the existence of $(z, \phi, \psi^1, \psi^2) \in L^1(\Omega) \times H^1(\Omega) \times \text{BV}^2$ such that $\phi_k \rightarrow \phi$ in $L^2(\Omega)$, $\phi_k \rightharpoonup \phi$ in $H^1(\Omega)$, $z_k \rightarrow z$ in $L^1(\Omega)$ and $\psi_k^j \rightarrow \psi^j$ in $L^1(\Omega)$. Moreover, z, ψ^1 and $\psi^2 \in \text{BV}$. See [21, Theorem 4, pp. 176].

From Lemma 5, we conclude that $(z, \phi, \psi^1, \psi^2)$ is an admissible quadruple. Moreover, from the weak lower semi-continuity of R (Lemma 8), together with the continuity of q (Lemma 3) and continuity of F (see the general assumption), we obtain

$$\begin{aligned} \inf \mathcal{G}_\alpha &= \lim_{k \rightarrow \infty} \mathcal{G}_\alpha(z_k, \phi_k, \psi_k^1, \psi_k^2) = \lim_{k \rightarrow \infty} \{ \|F(q(z_k, \psi_k^1, \psi_k^2)) - y^\delta\|_Y^2 + \alpha R(z_k, \phi_k, \psi_k^1, \psi_k^2) \} \\ &\geq \|F(q(z, \psi^1, \psi^2)) - y^\delta\|_Y^2 + \alpha R(z, \phi, \psi^1, \psi^2) = \mathcal{G}_\alpha(z, \phi, \psi^1, \psi^2), \end{aligned} \quad (15)$$

proving that $(z, \phi, \psi^1, \psi^2)$ minimizes $\mathcal{G}_\alpha$. $\square$

In that follows, we shall denote a minimizer of $\mathcal{G}_\alpha$ by $(z_\alpha, \phi_\alpha, \psi_\alpha^1, \psi_\alpha^2)$. In particular the functional $\hat{\mathcal{G}}_\alpha$ in (22) attain a generalized minimizer in the sense of Definition 1. In the next theorem, we summarize some convergence results for the regularized minimizers. These results are based on the existence of a generalized *minimum norm solutions*.

Definition 2. An admissible quadruple $(z^\dagger, \phi^\dagger, \psi^{1,\dagger}, \psi^{2,\dagger})$ is called a *R-minimizing solution* if satisfies

- (i) $F(q(z^\dagger, \psi^{1,\dagger}, \psi^{2,\dagger})) = y$,
- (ii) $R(z^\dagger, \phi^\dagger, \psi^{1,\dagger}, \psi^{2,\dagger}) = ms := \inf \{ R(z, \phi, \psi^1, \psi^2) ; (z, \phi, \psi^1, \psi^2) \text{ is an admissible quadruple and } F(q(z, \psi^1, \psi^2)) = y \}$.

Theorem 10 (*R-minimizing solutions*). Under the general assumptions of this paper there exists a *R-minimizing solution*.

Proof. From the general assumption on this paper and Remark 3, we conclude that the set of admissible quadruple satisfying $F(q(z, \psi^1, \psi^2)) = y$ is not empty. Thus, ms in (ii) is finite and there exists a sequence $\{(z_k, \phi_k, \psi_k^1, \psi_k^2)\}_{k \in \mathbb{N}}$ of admissible quadruple satisfying

$$F(q(z_k, \psi_k^1, \psi_k^2)) = y \quad \text{and} \quad R(z_k, \phi_k, \psi_k^1, \psi_k^2) \rightarrow ms < \infty.$$

Now, from the definition of R , it follows that the sequences $\{\phi_k\}_{k \in \mathbb{N}}$, $\{z_k\}_{k \in \mathbb{N}}$ and $\{\psi_k^j\}_{k \in \mathbb{N}}^{j=1,2}$ are uniformly bounded in $H^1(\Omega)$ and $\text{BV}(\Omega)$, respectively. Then, from the Sobolev Compact Embedding Theorem [2, 21], we have (up to subsequences) that

$$\phi_k \rightarrow \phi^\dagger \text{ in } L^2(\Omega), \quad z_k \rightarrow z^\dagger \text{ in } L^1(\Omega) \quad \text{and} \quad \psi_k^j \rightarrow \psi^{j,\dagger} \text{ in } L^1(\Omega), \quad j = 1, 2.$$

Lemma 5 implies that $(z^\dagger, \phi^\dagger, \psi^{1,\dagger}, \psi^{2,\dagger})$ is an admissible quadruple. Since R is weakly lower semi-continuous (cf. Lemma 8), it follows

$$ms = \liminf_{k \rightarrow \infty} R(z_k, \phi_k, \psi_k^1, \psi_k^2) \geq R(z^\dagger, \phi^\dagger, \psi^{1,\dagger}, \psi^{2,\dagger}).$$

Moreover, we conclude from Lemma 3 that

$$q(z^\dagger, \psi^{1,\dagger}, \psi^{2,\dagger}) = \lim_{k \rightarrow \infty} q(z_k, \psi_k^1, \psi_k^2) \quad \text{and} \quad F(q(z^\dagger, \psi^{1,\dagger}, \psi^{2,\dagger})) = \lim_{k \rightarrow \infty} F(q(z_k, \psi_k^1, \psi_k^2)) = y.$$

Thus, $(z^\dagger, \phi^\dagger, \psi^{1,\dagger}, \psi^{2,\dagger})$ is a *R-minimizing solution*. $\square$

Using classical techniques from the analysis of Tikhonov regularization methods (see [20, 19]), we present below the main convergence and stability theorems of this paper. The arguments in the proof are somewhat different of that presented in [14, 13]. But, for sake of completeness, we present the proof.

Theorem 11 (Convergence for exact data). *Assume that we have exact data, i.e. $y^\delta = y$. For every $\alpha > 0$ let $(z_\alpha, \phi_\alpha, \psi_\alpha^1, \psi_\alpha^2)$ denote a minimizer of $\mathcal{G}_\alpha$ on the set of admissible quadruples. Then, for every sequence of positive numbers $\{\alpha_k\}_{k \in \mathbb{N}}$ converging to zero there exists a subsequence, denoted again by $\{\alpha_k\}_{k \in \mathbb{N}}$, such that $(z_{\alpha_k}, \phi_{\alpha_k}, \psi_{\alpha_k}^1, \psi_{\alpha_k}^2)$ is strongly convergent in $L^1(\Omega) \times L^2(\Omega) \times (L^1(\Omega))^2$. Moreover, the limit is a solution of (1).*

Proof. Let $(z^\dagger, \phi^\dagger, \psi^{1,\dagger}, \psi^{2,\dagger})$ be a R -minimizing solution of (1) – its existence is guaranteed by Theorem 10. Let $\{\alpha_k\}_{k \in \mathbb{N}}$ be a sequence of positive numbers converging to zero. For each $k \in \mathbb{N}$, denote $(z_k, \phi_k, \psi_k^1, \psi_k^2) := (z_{\alpha_k}, \phi_{\alpha_k}, \psi_{\alpha_k}^1, \psi_{\alpha_k}^2)$ be a minimizer of G_{α_k} . Then, for each $k \in \mathbb{N}$, we have

$$G_{\alpha_k}(z_k, \phi_k, \psi_k^1, \psi_k^2) \leq \|F(q(z^\dagger, \psi^{1,\dagger}, \psi^{2,\dagger})) - y\| + \alpha_k R(z^\dagger, \phi^\dagger, \psi^{1,\dagger}, \psi^{2,\dagger}) = \alpha_k R(z^\dagger, \phi^\dagger, \psi^{1,\dagger}, \psi^{2,\dagger}). \quad (16)$$

Since $\alpha_k R(z_k, \phi_k, \psi_k^1, \psi_k^2) \leq G_{\alpha_k}(z_k, \phi_k, \psi_k^1, \psi_k^2)$, it follows from (16) that

$$R(z_k, \phi_k, \psi_k^1, \psi_k^2) \leq R(z^\dagger, \phi^\dagger, \psi^{1,\dagger}, \psi^{2,\dagger}) < \infty. \quad (17)$$

Moreover, from the assumption on the sequence $\{\alpha_k\}$, it follows that

$$\lim_{k \rightarrow \infty} \alpha_k R(z^\dagger, \phi^\dagger, \psi^{1,\dagger}, \psi^{2,\dagger}) = 0. \quad (18)$$

From (17) and Lemma 7, we conclude that sequences $\{\phi_k\}$, $\{z_k\}$ and $\{\psi_k^j\}$ are bounded in $H^1(\Omega)$ and BV, respectively, for $j = 1, 2$. Using an argument of extraction of diagonal subsequences (see proof of Lemma 5), we can guarantee the existence of an admissible quadruple $(\tilde{z}, \tilde{\phi}, \tilde{\psi}^1, \tilde{\psi}^2)$ such that

$$(z_k, \phi_k, \psi_k^1, \psi_k^2) \rightarrow (\tilde{z}, \tilde{\phi}, \tilde{\psi}^1, \tilde{\psi}^2) \text{ in } L^1(\Omega) \times L^2(\Omega) \times (L^1(\Omega))^2.$$

Now, from Lemma 3 (i), it follows that $q(\tilde{z}, \tilde{\psi}^1, \tilde{\psi}^2) = \lim_{k \rightarrow \infty} q(z_k, \psi_k^1, \psi_k^2)$ in $L^1(\Omega)$. Using the continuity of the operator F together with (16) and (18), we conclude that

$$y = \lim_{k \rightarrow \infty} F(q(z_k, \psi_k^1, \psi_k^2)) = F(q(\tilde{z}, \tilde{\psi}^1, \tilde{\psi}^2)).$$

On the other hand, from the lower semi-continuity of R and (17) it follows that

$$R(\tilde{z}, \tilde{\phi}, \tilde{\psi}^1, \tilde{\psi}^2) \leq \liminf_{k \rightarrow \infty} R(z_k, \phi_k, \psi_k^1, \psi_k^2) \leq \limsup_{k \rightarrow \infty} R(z_k, \phi_k, \psi_k^1, \psi_k^2) \leq R(z^\dagger, \phi^\dagger, \psi^{1,\dagger}, \psi^{2,\dagger}),$$

concluding the proof. $\square$

Theorem 12 (Stability). *Let $\alpha = \alpha(\delta)$ be a function satisfying $\lim_{\delta \rightarrow 0} \alpha(\delta) = 0$ and $\lim_{\delta \rightarrow 0} \delta^2 \alpha(\delta)^{-1} = 0$. Moreover, let $\{\delta_k\}_{k \in \mathbb{N}}$ be a sequence of positive numbers converging to zero and $y^{\delta_k} \in Y$ be corresponding noisy data satisfying (2). Then, there exist a subsequence, denoted again by $\{\delta_k\}$, and a sequence $\{\alpha_k := \alpha(\delta_k)\}$ such that $(z_{\alpha_k}, \phi_{\alpha_k}, \psi_{\alpha_k}^1, \psi_{\alpha_k}^2)$ converges in $L^1(\Omega) \times L^2(\Omega) \times (L^1(\Omega))^2$ to solution of (1).*

Proof. Let $(z^\dagger, \phi^\dagger, \psi^{1,\dagger}, \psi^{2,\dagger})$ be a R -minimizer solution of (1) (such existence is guaranteed by Theorem 10). For each $k \in \mathbb{N}$, let $(z_k, \phi_k, \psi_k^1, \psi_k^2) := (z_{\alpha(\delta_k)}, \phi_{\alpha(\delta_k)}, \psi_{\alpha(\delta_k)}^1, \psi_{\alpha(\delta_k)}^2)$ be a minimizer of $G_{\alpha(\delta_k)}$. Then, for each $k \in \mathbb{N}$ we have

$$\begin{aligned} G_{\alpha_k}(z_k, \phi_k, \psi_k^1, \psi_k^2) &\leq \|F(q(z^\dagger, \psi^{1,\dagger}, \psi^{2,\dagger})) - y^{\delta_k}\|_Y^2 + \alpha(\delta_k) R(z^\dagger, \phi^\dagger, \psi^{1,\dagger}, \psi^{2,\dagger}) \\ &\leq \delta_k^2 + \alpha(\delta_k) R(z^\dagger, \phi^\dagger, \psi^{1,\dagger}, \psi^{2,\dagger}). \end{aligned} \quad (19)$$

From (19) and the definition of G_{α_k} , it follows that

$$R(z_k, \phi_k, \psi_k^1, \psi_k^2) \leq \frac{\delta_k^2}{\alpha(\delta_k)} + R(z^\dagger, \phi^\dagger, \psi^{1,\dagger}, \psi^{2,\dagger}). \quad (20)$$

Taking the limit as $k \rightarrow \infty$ in (20), it follows from theorem assumptions on $\alpha(\delta_k)$, that

$$\lim_{k \rightarrow \infty} \left\| F(q(z_k, \psi_k^1, \psi_k^2)) - y^{\delta_k} \right\| \leq \lim_{k \rightarrow \infty} \left(\delta_k^2 + \alpha(\delta_k) R(z^\dagger, \phi^\dagger, \psi^{1,\dagger}, \psi^{2,\dagger}) \right) = 0,$$

and

$$\limsup_{k \rightarrow \infty} R(z_k, \phi_k, \psi_k^1, \psi_k^2) \leq R(z^\dagger, \phi^\dagger, \psi^{1,\dagger}, \psi^{2,\dagger}). \quad (21)$$

With the same arguments as in the proof of Theorem 11, we conclude that, at least a subsequence that we denote again by $(z_k, \phi_k, \psi_k^1, \psi_k^2)$, converge in $L^1(\Omega) \times L^2(\Omega) \times (L^1(\Omega))^2$ to some admissible quadruple $(z, \phi, \psi^1, \psi^2)$. Moreover, by taking the limit as $k \rightarrow \infty$ in (19), it follows from the assumption on F and Lemma 3 that

$$F(q(z, \phi, \psi^1, \psi^2)) = \lim_{k \rightarrow \infty} F(q(z_k, \psi_k^1, \psi_k^2)) = y.$$

□

The functional $\mathcal{G}_\alpha$ defined in (7) is not easy to handled numerically, i.e., we are not able to derive a suitable optimality condition to the minimizers of $\mathcal{G}_\alpha$. In the next section, we work in sight to surpass such difficulty.

5 Numerical Solution

In this section, we introduce a functional which can be handled numerically, and whose minimizers are 'near' to the minimizers of $\mathcal{G}_\alpha$. Let $\mathcal{G}_{\varepsilon, \alpha}$ be the functional defined by

$$\mathcal{G}_{\varepsilon, \alpha}(\phi, \psi^1, \psi^2) := \|F(P_\varepsilon(\phi, \psi^1, \psi^2)) - y^\delta\|_Y^2 + \alpha(\beta_1 |H_\varepsilon(\phi)|_{\text{BV}} + \beta_2 \|\phi - \phi_0\|_{H^1}^2 + \beta_3 \sum_{j=1}^2 |\psi^j - \psi_0^j|_{\text{BV}}), \quad (22)$$

where $P_\varepsilon(\phi, \psi^1, \psi^2) := q(H_\varepsilon(\phi), \psi^1, \psi^2)$ is defined in (6). The functional $\mathcal{G}_{\varepsilon, \alpha}$ is well-posed as the following lemma shows:

Lemma 13. *Given positive constants $\alpha, \varepsilon, \beta_j$ as in the general assumption of this article, $\phi_0 \in H^1(\Omega)$ and $\psi_0^j \in \mathbb{B}$, $j = 1, 2$. Then, the functional $\mathcal{G}_{\varepsilon, \alpha}$ in (22) attains a minimizer on $H^1(\Omega) \times (\text{BV})^2$.*

Proof. Since, $\inf\{\mathcal{G}_{\varepsilon, \alpha}(\phi, \psi^1, \psi^2) : (\phi, \psi^1, \psi^2) \in H^1(\Omega) \times (\text{BV})^2\} \leq \mathcal{G}_{\varepsilon, \alpha}(0, 0, 0) < \infty$, there exists a minimizing sequence $\{(\phi_k, \psi_k^1, \psi_k^2)\}$ in $H^1(\Omega) \times \mathbb{B}^2$ satisfying

$$\lim_{k \rightarrow \infty} \mathcal{G}_{\varepsilon, \alpha}(\phi_k, \psi_k^1, \psi_k^2) = \inf\{\mathcal{G}_{\varepsilon, \alpha}(\phi, \psi^1, \psi^2) : (\phi, \psi^1, \psi^2) \in H^1(\Omega) \times \mathbb{B}^2\}.$$

Then, for fixed $\alpha > 0$, the definition of $\mathcal{G}_{\varepsilon, \alpha}$ in (22) implies that the sequences $\{\phi_k\}$ and $\{\psi_k^j\}_{j=1,2}$ are bounded in $H^1(\Omega)$ and $(\text{BV})^2$, respectively. Therefore, from Banach-Alaoglu-Bourbaki Theorem [42] $\phi_k \rightharpoonup \phi$ in $H^1(\Omega)$ and from [21, Theorem 4 pg. 176], $\psi_k^j \rightarrow \psi^j$ in $L^1(\Omega)$, $j = 1, 2$. Now, a similar argument as in Lemma 4 implies that $\psi^j \in \mathbb{B}$, for $j = 1, 2$. Moreover, by the weak lower semi-continuity of the H^1 -norm [42] and $|\cdot|_{\text{BV}}$ measure (see [21, Theorem 1 pg. 172]), it follows that

$$\|\phi - \phi_0\|_{H^1}^2 \leq \liminf_{k \rightarrow \infty} \|\phi_k - \phi_0\|_{H^1}^2 \quad \text{and} \quad |\psi^j - \psi_0^j|_{\text{BV}} \leq \liminf_{k \rightarrow \infty} |\psi_k^j - \psi_0^j|_{\text{BV}}.$$

The compact embedding of $H^1(\Omega)$ into $L^2(\Omega)$ [2] implies in the existence of a subsequence of $\{\phi_k\}$, (that we denote with the same index) such that $\phi_k \rightarrow \phi$ in $L^2(\Omega)$. Follows from Lemma 3 and [21,

Theorem 1, pg 172] that $|H_\varepsilon(\phi)|_{\text{BV}} \leq \liminf_{k \rightarrow \infty} |H_\varepsilon(\phi_k)|_{\text{BV}}$. Hence, from continuity of F in L^1 , continuity of q (see Lemma 3), together with the estimates above, we conclude that

$$\begin{aligned} \mathcal{G}_{\varepsilon,\alpha}(\phi, \psi^1, \psi^2) &\leq \lim_{k \rightarrow \infty} \|F(P_\varepsilon(\phi_k, \psi_k^1, \psi_k^2)) - y^\delta\|_Y^2 \\ &\quad + \alpha \left(\beta_1 \liminf_{k \rightarrow \infty} |H_\varepsilon(\phi_k)|_{\text{BV}} + \beta_2 \liminf_{k \rightarrow \infty} \|\phi_k - \phi_0\|_{H^1(\Omega)}^2 + \beta_3 \liminf_{k \rightarrow \infty} \sum_{j=1}^2 |\psi_k^j - \psi_0^j|_{\text{BV}} \right) \\ &\leq \liminf_{k \rightarrow \infty} \mathcal{G}_{\varepsilon,\alpha}(\phi_k, \psi_k^1, \psi_k^2) = \inf \mathcal{G}_{\varepsilon,\alpha}, \end{aligned}$$

Therefore, (ϕ, ψ^1, ψ^2) is a minimizer of $\mathcal{G}_{\varepsilon,\alpha}$. $\square$

In the sequel, we prove that, when $\varepsilon \rightarrow 0$, the minimizers of $\mathcal{G}_{\varepsilon,\alpha}$ approximate a minimizer of the functional $\mathcal{G}_\alpha$. Hence, numerically, the minimizer of $\mathcal{G}_{\varepsilon,\alpha}$ can be used as a suitable approximation for the minimizers of $\mathcal{G}_\alpha$.

Theorem 14. *Let α and β_j be given as in the general assumption of this article. For each $\varepsilon > 0$, denote by $(\phi_{\varepsilon,\alpha}, \psi_{\varepsilon,\alpha}^1, \psi_{\varepsilon,\alpha}^2)$ a minimizer of $\mathcal{G}_{\varepsilon,\alpha}$ (that there exist from Lemma 13). Then, there exists a sequence of positive numbers $\varepsilon_k \rightarrow 0$ such that $(H_{\varepsilon_k}(\phi_{\varepsilon_k,\alpha}), \phi_{\varepsilon_k,\alpha}, \psi_{\varepsilon_k,\alpha}^1, \psi_{\varepsilon_k,\alpha}^2)$ converges strongly in $L^1(\Omega) \times L^2(\Omega) \times (L^1(\Omega))^2$ and the limit minimizes $\mathcal{G}_\alpha$ on the set of admissible quadruples.*

Proof. Let $(z_\alpha, \phi_\alpha, \psi_\alpha^1, \psi_\alpha^2)$ be a minimizer of the functional $\mathcal{G}_\alpha$ on the set of admissible quadruples (cf. Theorem 9). From Definition 1, there exists a sequence $\{\varepsilon_k\}$ of positive numbers converging to zero and corresponding sequences $\{\phi_k\}$ in $H^1(\Omega)$ satisfying $\phi_k \rightarrow \phi_\alpha$ in $L^2(\Omega)$, $H_{\varepsilon_k}(\phi_k) \rightarrow z_\alpha$ in $L^1(\Omega)$ and, finally, sequences $\{\psi_k^j\}$ in $\text{BV} \times C_c^\infty(\Omega)$ such that $|\psi_k^j|_{\text{BV}} \rightarrow |\psi^j|_{\text{BV}}$. Moreover, we can further assume (see Lemma 6) that

$$R(z_\alpha, \phi_\alpha, \psi_\alpha^1, \psi_\alpha^2) = \lim_{k \rightarrow \infty} (\beta_1 |H_{\varepsilon_k}(\phi_k)|_{\text{BV}} + \beta_2 \|\phi_k - \phi_0\|_{H^1(\Omega)}^2 + \beta_3 \sum_{j=1}^2 |\psi_k^j - \psi_0^j|_{\text{BV}}).$$

Let $(\phi_{\varepsilon_k}, \psi_{\varepsilon_k}^1, \psi_{\varepsilon_k}^2)$ be a minimizer of $\mathcal{G}_{\varepsilon_k,\alpha}$. Hence, $(\phi_{\varepsilon_k}, \psi_{\varepsilon_k}^1, \psi_{\varepsilon_k}^2)$ belongs to $H^1(\Omega) \times \mathbb{B}^2$ (see Lemma 13). The sequences $\{H_{\varepsilon_k}(\phi_{\varepsilon_k})\}, \{\phi_{\varepsilon_k}\}$ and $\{\psi_{\varepsilon_k}^j\}$ are uniformly bounded in $\text{BV}(\Omega)$, $H^1(\Omega)$ and $\text{BV}(\Omega)$, for $j = 1, 2$, respectively. From compact embedding (see Theorems [2] and [21, Theorem 4 pg. 176]), there exist convergent subsequences whose limits are denoted by $\tilde{z}$, $\tilde{\phi}$ and $\tilde{\psi}^j$ belong to $\text{BV}(\Omega), H^1(\Omega)$ and $\text{BV}(\Omega)$, for $j = 1, 2$, respectively.

Summarizing, we have $\phi_{\varepsilon_k} \rightarrow \tilde{\phi}$ in $L^2(\Omega)$, $H_{\varepsilon_k}(\phi_{\varepsilon_k}) \rightarrow \tilde{z}$ in $L^1(\Omega)$, and $\psi_{\varepsilon_k}^j \rightarrow \tilde{\psi}^j$ in $L^1(\Omega)$, $j = 1, 2$. Thus, $(\tilde{z}, \tilde{\phi}, \tilde{\psi}^1, \tilde{\psi}^2) \in L^1(\Omega) \times H^1(\Omega) \times L^1(\Omega)$ is an admissible quadruple (cf. Lemma 5).

From the definition of R , Lemma 3 and the continuity of F , it follows that

$$\begin{aligned} \|F(q(\tilde{z}, \tilde{\psi}^1, \tilde{\psi}^2)) - y^\delta\|_Y^2 &= \lim_{k \rightarrow \infty} \|F(P_{\varepsilon_k}(\phi_{\varepsilon_k}, \psi_{\varepsilon_k}^1, \psi_{\varepsilon_k}^2)) - y^\delta\|_Y^2, \\ R(\tilde{z}, \tilde{\phi}, \tilde{\psi}^1, \tilde{\psi}^2) &\leq \liminf_{k \rightarrow \infty} (\beta_1 |H_{\varepsilon_k}(\phi_{\varepsilon_k})|_{\text{BV}} + \beta_2 \|\phi_{\varepsilon_k} - \phi_0\|_{H^1(\Omega)}^2 + \beta_3 \sum_{j=1}^2 |\psi_{\varepsilon_k}^j - \psi_0^j|_{\text{BV}}). \end{aligned}$$

Therefore,

$$\begin{aligned} \mathcal{G}_\alpha(\tilde{z}, \tilde{\phi}, \tilde{\psi}^1, \tilde{\psi}^2) &= \|F(q(\tilde{z}, \tilde{\psi}^1, \tilde{\psi}^2)) - y^\delta\|_Y^2 + \alpha R(\tilde{z}, \tilde{\phi}, \tilde{\psi}^1, \tilde{\psi}^2) \\ &\leq \liminf_{k \rightarrow \infty} \mathcal{G}_{\varepsilon_k,\alpha}(\phi_{\varepsilon_k}, \psi_{\varepsilon_k}^1, \psi_{\varepsilon_k}^2) \leq \liminf_{k \rightarrow \infty} \mathcal{G}_{\varepsilon_k,\alpha}(\phi_k, \psi_k^1, \psi_k^2) \\ &\leq \limsup_{k \rightarrow \infty} \|F(P_{\varepsilon_k}(\phi_k, \psi_k^1, \psi_k^2)) - y^\delta\|_Y^2 \\ &\quad + \alpha \limsup_{k \rightarrow \infty} (\beta_1 |H_{\varepsilon_k}(\phi_k)|_{\text{BV}} + \beta_2 \|\phi_k - \phi_0\|_{H^1(\Omega)}^2 + \beta_3 \sum_{j=1}^2 |\psi_k^j - \psi_0^j|_{\text{BV}}) \\ &= \|F(q(z_\alpha, \psi_\alpha^1, \psi_\alpha^2)) - y^\delta\|_Y^2 + \alpha R(z_\alpha, \phi_\alpha, \psi_\alpha^1, \psi_\alpha^2) = \mathcal{G}_\alpha(z_\alpha, \phi_\alpha, \psi_\alpha^1, \psi_\alpha^2), \end{aligned}$$

characterizing $(\tilde{z}, \tilde{\phi}, \tilde{\psi}^1, \tilde{\psi}^2)$ as a minimizer of $\mathcal{G}_\alpha$. $\square$

5.1 Optimality Conditions for the Stabilized Functional

For numerical purposes it is convenient to derive first order optimality conditions for minimizers of the functional $\mathcal{G}_\alpha$. Since P is a discontinuous operator, it is not possible. However, thanks to the Theorem 12, the minimizers of the stabilized functionals $\mathcal{G}_{\varepsilon,\alpha}$ can be used for approximate minimizers of the functional $\mathcal{G}_\alpha$. Therefore, we consider $\mathcal{G}_{\varepsilon,\alpha}$ in (22), with Y a Hilbert space, and we look for the Gâteaux directional derivatives with respect to ϕ and the unknown ψ^j for $j = 1, 2$.

Since $H'_\varepsilon(\phi)$ is self-adjoint¹, we can write the optimality conditions for the functional $\mathcal{G}_{\varepsilon,\alpha}$ in the form of the system

$$\alpha(\Delta - I)(\phi - \phi_0) = L_{\varepsilon,\alpha,\beta}(\phi, \psi^1, \psi^2), \quad \text{in } \Omega \quad (23a)$$

$$(\phi - \phi_0) \cdot \nu = 0, \quad \text{at } \partial\Omega \quad (23b)$$

$$\alpha \nabla \cdot [\nabla(\psi^j - \psi_0^j)/|\nabla(\psi^j - \psi_0^j)|] = L_{\varepsilon,\alpha,\beta}^j(\phi, \psi^1, \psi^2), \quad j = 1, 2. \quad (23c)$$

Here $\nu(x)$ represents the external unit normal quadruple at $x \in \partial\Omega$, and

$$\begin{aligned} L_{\varepsilon,\alpha,\beta}(\phi, \psi^1, \psi^2) &= (\psi^1 - \psi^2)\beta_2^{-1}H'_\varepsilon(\phi)^*F'(P_\varepsilon(\phi, \psi^1, \psi^2))^*(F(P_\varepsilon(\phi, \psi^1, \psi^2)) - y^\delta) \\ &\quad - \beta_1(2\beta_2)^{-1}H'_\varepsilon(\phi) \nabla \cdot [\nabla H_\varepsilon(\phi)/|\nabla H_\varepsilon(\phi)|], \end{aligned} \quad (24a)$$

$$L_{\varepsilon,\alpha,\beta}^1(\phi, \psi^1, \psi^2) = (2\beta_3)^{-1}(F'(P_\varepsilon(\phi, \psi^1, \psi^2))H_\varepsilon(\phi))^*(F(P_\varepsilon(\phi, \psi^1, \psi^2)) - y^\delta) \quad (24b)$$

$$L_{\varepsilon,\alpha,\beta}^2(\phi, \psi^1, \psi^2) = (2\beta_3)^{-1}(F'(P_\varepsilon(\phi, \psi^1, \psi^2))(1 - H_\varepsilon(\phi)))^*(F(P_\varepsilon(\phi, \psi^1, \psi^2)) - y^\delta). \quad (24c)$$

It is worth noticing that the derivation of (23) is purely formal, since the BV seminorm is not differentiable. Moreover the terms $|\nabla H_\varepsilon(\phi)|$ and $|\nabla(\psi^j - \psi_0^j)|$ appearing in the denominators of (23) and (24), respectively.

In Section 6, system (23) and (24) is used as starting point for the derivation of a level set type method.

6 Inverse Elliptic Problems

In this section, we discuss the proposed level set approach and their application in some physical problems model by elliptic PDE's. We also discuss briefly the numerical implementations of the iterative method based on the level set approach. We remark that the iterative algorithm derived by the level set approach need a early stoop criteria [19], in the case of noise data.

6.1 The Inverse Potential Problem

In this subsection, we apply the level set regularization framework developed before in this article in an inverse potential problem, similar to ones considered in [38, 39, 13, 14, 22, 36, 23, 37].

The forward problem consists of solving on a given Lipschitz domain $\Omega \subset \mathbb{R}^n$, for a given source function $u \in L^2(\Omega)$ and a boundary function $g \in L^2(\partial\Omega)$, the Poisson boundary value problem

$$-\nabla \cdot (\sigma \nabla w) = u, \text{ in } \Omega, \quad \gamma_1 w + \gamma_2 w_\nu = g \text{ on } \partial\Omega. \quad (25)$$

In (25), ν represent the outer normal vector to $\partial\Omega$, σ is a known sufficient smooth function. Note that, depending of $\gamma_1, \gamma_2 \in \{0, 1\}$, we have Dirichlet, Neumann or Robin boundary condition. In this paper, we only consider the case of Dirichlet boundary condition, that corresponds to $\gamma_1 = 1$ and $\gamma_2 = 0$ in (25). It is well known that there exists a unique solution $w \in H^1(\Omega)$ of (25) with $w - g \in H_0^1(\Omega)$, [12].

These simple version of the elliptic problem (25) appears in many relevant applications including: inverse potential problems [23], inverse gravimetry [24] and EMG [39, 38].

¹Note that $H'_\varepsilon(t) = \begin{cases} \frac{1}{\varepsilon} & t \in (-\varepsilon, 0) \\ 0 & \text{other else.} \end{cases}$

Assuming homogeneous Dirichlet boundary condition in (25), the problem can be modeled by the operator equation

$$\begin{aligned} F_1 &: L^2(\Omega) \rightarrow L^2(\partial\Omega) \\ u &\mapsto F_1(u) := w_\nu|_{\partial\Omega}. \end{aligned} \quad (26)$$

The corresponding inverse problem is, in general, called *inverse potential problem*, which consists of recovering an L^2 -function u , from measurements of the Cauchy data of its corresponding potential on the boundary of Ω .

Using this notation, the inverse potential problem can be written in the abbreviated form $F_1(u) = y^\delta$, where the available noisy data $y^\delta \in L^2(\partial\Omega)$ have the same meaning as in (2). It is worth noticing that this inverse problem has, in general, non unique solution [23]. Sufficient conditions for identifiability are given in [24]. Moreover, in this case, the operator F_1 is linear. However, the inverse potential problem is well known to be exponentially ill-posed [24]. Therefore, the solution call for a regularization strategy [19, 23, 24].

Given the non-uniqueness of the inverse potential problem and the limitation of our approach, we restrict our attention to solve the inverse problem (26) in $D(F)$, i.e., we assume that the unknown parameter $u \in D(F)$, as defined in Section 3. For this class of parameters no unique identifiability result is known and we restrict our attention to minimum-norm solutions [19]. The following lemma implies that the operator F_1 satisfies the Assumption (A2).

Lemma 15. *The operator $F_1 : D(F) \subset L^1(\Omega) \longrightarrow L^2(\partial\Omega)$ is continuous with the respect to the $L^1(\Omega)$ topology.*

Proof. It is well known from the elliptic regularity theory [12] that $\|w\|_{H^1(\Omega)} \leq c_1\|u\|_{L^2(\Omega)}$. Let $u_n, u_0 \in D(F)$ and w_n, w_0 the respective solution of (25). Then, the linearity and continuity of the trace operator from $H^1(\Omega)$ to $L^2(\partial\Omega)$ [12], we have that

$$\|F_1(u_n) - F_1(u_0)\|_{L^2(\partial\Omega)} \leq C\|w_n - w_0\|_{H^1(\Omega)} \leq \tilde{C}\|u_n - u_0\|_{L^2(\Omega)}.$$

From Lemma 2, we conclude that F_1 is sequentially continuous on the $L^1(\Omega)$ topology. Since $L^1(\Omega)$ is a metrizable spaces [42], the proof is complete. $\square$

6.1.1 A level set algorithm for the inverse potential problem

The iterative algorithm based on the level set approach proposed in this article is an explicit iterative method derived from the optimality conditions for the Tikhonov functional $\mathcal{G}_{\varepsilon, \alpha}$ in (23) and (24). To the inverse potential problem for Dirichlet boundary condition ($\gamma_1 = 1$ and $\gamma_2 = 0$) the algorithm reads as:

Each step of this iterative method consists of three parts (see Table 1): 1) The residual $r_k \in L^2(\partial\Omega)$ of the iterate (ϕ_k, ψ_k^j) is evaluated (this requires solving one elliptic BVP of Dirichlet type); 2) The L^2 -solution h_k of the adjoint problem for the residual is evaluated (this corresponds to solving one elliptic BVP of Dirichlet type); 3) The update $\delta\phi_k$ for the level-set function and the updates $\delta\psi_k^j$ for the level values are evaluated (this corresponds to multiplying two functions).

In [37], a level set method was proposed, where the iteration is based on an inexact Newton type method with the inner iteration is implemented using the conjugate gradient method. Moreover, the regularization parameter $\alpha > 0$ is kept fixed. In the above algorithm, we define $\delta t = 1/\alpha$ as time increment in order to derive an evolution equation for the levelset function. Therefore, we are looking for a fixed point equation related to the system of optimality conditions for the Tikhonov functional. Here the iteration is based on a gradient type method as in [13].

Given σ and g ;

1. Evaluate the residual $r_k := F_1(P_\varepsilon(\phi_k, \psi_k^1, \psi_k^2)) - y^\delta = (w_k)_\nu|_{\partial\Omega} - y^\delta$, where w_k solves $-\nabla \cdot (\sigma \nabla w_k) = P_\varepsilon(\phi_k, \psi_k^1, \psi_k^2)$, in Ω ; $w_k = g$, at $\partial\Omega$.
2. Evaluate $h_k := F_1'(P_\varepsilon(\phi_k, \psi_k^1, \psi_k^2))^*(r_k) \in L^2(\Omega)$, solving $\Delta h_k = 0$, in Ω ; $h_k = r_k$, at $\partial\Omega$.
3. Calculate $\delta\phi_k := L_{\varepsilon, \alpha, \beta}(\phi_k, \psi_k^1, \psi_k^2)$ and $\delta\psi_k^j := L_{\varepsilon, \alpha, \beta}^j(\phi_k, \psi_k^1, \psi_k^2)$, as in (24).
4. Update the level set function ϕ_k and the level values ψ_k^j , $j = 1, 2$:

$$\phi_{k+1} = \phi_k + \frac{1}{\alpha} \delta\phi_k, \quad \psi_{k+1}^j = \psi_k^j + \frac{1}{\alpha} \delta\psi_k^j.$$

Table 1: Iterative algorithm based on the level set approach for the inverse potential problem.

6.2 The Inverse Problem in Nonlinear Electromagnetism

Many interesting physical problems are model by quasi-linear elliptic equations. One such problem that is interesting in applications is the identification of inhomogeneity inside nonlinear magnetic materials form indirect or local measurements. Electromagnetic non-destructive tests aim to localize cracks or inhomogeneities in the steel production process where the impurities can be described by a piecewise smooth function, [8, 9, 5, 11].

In this section, we assume that $D \subset \subset \Omega$ is measurable and

$$u = \begin{cases} \psi_1, & x \in D, \\ \psi_2, & x \in \Omega \setminus D, \end{cases} \quad (27)$$

with $\psi_1, \psi_2 \in \mathbb{B}$ with $m > 0$.

The forward problem consists of solving on a given Lipschitz domain $\Omega \subset \mathbb{R}^n$, for a given source function $f \in H^{-1}(\Omega)$, a boundary function $g \in H^{1/2}(\partial\Omega)$, the Poisson boundary value problem

$$-\nabla \cdot (u \nabla w) = f, \text{ in } \Omega, \quad w = g \text{ on } \partial\Omega. \quad (28)$$

It is well known that there exists a unique solution $w \in H^1(\Omega)$ such that $w - g \in H_0^1(\Omega)$ for the problem (28), [12].

Assuming that, during the production process, the workpiece is contaminated by impurities and that such impurities are described by piecewise smooth function, the inverse electromagnetic problem consist in the identification and the localization of the inhomogeneities as well as the function values of the impurities. The localization of the support of the inhomogeneities and the tabulation of his values can indicate possible sources of contamination in the magnetic material. Note that, if ψ_1 and ψ_2 are constant, then (28) describe the EIT problem [36, 24, 6, 7].

The inverse problem that we are interesting here consists in the identification of the support (shape) and the function values of ψ^1, ψ^2 of the coefficient function $u(x)$ defined in (27), where the voltage potential g is chosen such that its corresponding the current measurement $h := (w)_\nu|_{\partial\Omega}$ are available as a set of continuous measurement in $\partial\Omega$. This problem is known in the literature as the inverse problem for the Dirichlet-to-Neumann operator [24].

With this framework, the problem can be model by the operator equation

$$\begin{aligned} F_2 : D(F) \subset L^1(\Omega) &\rightarrow H^{1/2}(\partial\Omega) \\ u &\mapsto F_2(u) := w|_{\partial\Omega}, \end{aligned} \quad (29)$$

where the potential profile $g = w|_{\partial\Omega} \in H^{1/2}(\Omega)$ is given.

In [11], a level set approach for solve an inverse problems of identification of inhomogeneities inside a nonlinear material form local measurements of the magnetic induction was investigate. The assumption in [11] is that part of the inhomogeneities are given by a crack localized inside the workpiece and that outside the crack region, magnetic conductivities are nonlinear and depends on the magnetic induction. In other words, that $\psi_1 = \mu_1$ and $\psi_2 = \mu_2(|\nabla w|^2)$, where μ_1 is the (constant) air conductivity and $\mu_2 = \mu_2(|\nabla w|^2)$ is a nonlinear conductivity of the workpiece material, whose values are assumed be known. A successful iterative algorithm and numerical experiment was presented. However, we remark that, in [11], the measurements and therefore the data are given in the whole Ω . Such measurements are not reasonable in applications. Moreover, the proposed level set algorithm is based on an optimality condition of a least square functional with $H^1(\Omega)$ -semi-norm regularization. However, there is no guarantee of existence of minimum for the proposed functional.

In the following lemma, we prove that the operator F_2 satisfies the Assumption (A2).

Remark 4. Note that the operator $F_2(u) = T_D(w)$, where T_D is the Dirichlet trace operator. Moreover, $T_D : H^1(\Omega) \rightarrow H^{1/2}(\partial\Omega)$ is linear and continuous [12], i.e., $\|T_D(w)\|_{H^{1/2}(\partial\Omega)} \leq c\|w\|_{H^1(\Omega)}$.

Lemma 16. Let the operator $F_2 : D(F) \subset L^1(\Omega) \rightarrow H^{1/2}(\partial\Omega)$ as defined in (29). Then, F_2 is continuous with the respect to the $L^1(\Omega)$ topology.

Proof. Let $u_n, u_0 \in D(F)$ and w_n, w_0 denoting the respective solution of (25). The linearity of equation (28) implies that $w_n - w_0 \in H_0^1(\Omega)$ satisfies

$$\nabla \cdot (u_n \nabla w_n) - \nabla \cdot (u_0 \nabla w_0) = 0, \quad (30)$$

with homogeneous boundary condition. Therefore, using the weak formulation for (30) we have

$$\int_{\Omega} (\nabla \cdot (u_n \nabla w_n) - \nabla \cdot (u_0 \nabla w_0)) \varphi dx = 0, \quad \forall \varphi \in H_0^1(\Omega).$$

In particular, the weak formulation holds true for $\varphi = w_n - w_0$. From the Green formula [12] and the assumption that $m > 0$ (that guarantee ellipticity of (28)), follows that

$$m\|\nabla w_n - \nabla w_0\|_{L^2(\Omega)}^2 \leq \int_{\Omega} u_n |\nabla w_n - \nabla w_0|^2 dx \leq \int_{\Omega} (u_n - u_0) |\nabla w_0| |\nabla w_n - \nabla w_0| dx. \quad (31)$$

From [28, Theorem 1], there exist $\varepsilon > 0$ (small enough) such that $w_0 \in W^{1,p}(\Omega)$ for $p = 2 + \varepsilon$. Using the Hölder inequality [12] with $1/p + 1/q = 1/2$ (note that $q > 2$ in the equation (31)), follows that

$$m\|\nabla w_n - \nabla w_0\|_{L^2(\Omega)}^2 \leq \|u_n - u_0\|_{L^q(\Omega)} \|\nabla w_0\|_{L^p(\Omega)} \|\nabla w_n - \nabla w_0\|_{L^2(\Omega)}. \quad (32)$$

Therefore, using the Poincaré inequality [12] and equation (32), we have

$$\|w_n - w_0\|_{H^1(\Omega)} \leq C \|u_n - u_0\|_{L^q(\Omega)},$$

where the constant C depends only of $m, \Omega, \|\nabla w_0\|$ and the Poincaré constant. Now, the assertion follows from Lemma 2 and Remark 4. $\square$

6.2.1 A level set algorithm for inverse problem in nonlinear electromagnetism

The explicit iterative algorithm based on the level set approach proposed in this article is based on the optimality conditions for the Tikhonov functional $\mathcal{G}_{\varepsilon, \alpha}$ in (23) and (24). Each iteration of this algorithm consists in the following steps: In the first step the residual vector $r \in L^2(\partial\Omega)$ corresponding to the iterate $(\phi_n, \psi_n^1, \psi_n^2)$ is evaluated. This requires the solution of one elliptic BVP's of Dirichlet type. In the second step the solutions $v \in H^1(\Omega)$ of the adjoint problems for the residual components r are evaluated. This corresponds to solving one elliptic BVP of Neumann type and to computing the inner-product $\nabla w \cdot \nabla v$ in $L^2(\Omega)$. Next, the computation of $L_{\varepsilon, \alpha, \beta}(\phi_n, \psi_n^1, \psi_n^2)$ and $L_{\varepsilon, \alpha, \beta}^j(\phi_n, \psi_n^1, \psi_n^2)$ as in (24). The four step is the updates of the level-set function $\delta\phi_n \in H^1(\Omega)$ and the level function values $\delta\psi_n^j \in \text{BV}(\Omega)$ by solve (23).

The algorithm is summarized in Table 2.

- | |
|--|
| <p>1. Evaluate the residual $r := F_2(P_\varepsilon(\phi_n, \psi_n^1, \psi_n^2)) - y^\delta = w _{\partial\Omega} - g^\delta$, where $w \in H^1(\Omega)$ solves</p> $\nabla \cdot (P_\varepsilon(\phi_n, \psi_n^1, \psi_n^2) \nabla w) = f, \text{ in } \Omega; \quad w = g, \text{ at } \partial\Omega.$ <p>2. Evaluate $F_2'(P_\varepsilon(\phi_n, \psi_n^1, \psi_n^2))^* r := \nabla w \cdot \nabla v \in L^2(\Omega)$, where w is the function computed in Step 1. and $v \in H^1(\Omega)$ solves</p> $\nabla \cdot (P_\varepsilon(\phi_n, \psi_n^1, \psi_n^2) \nabla v) = 0 \text{ in } \Omega; \quad v_\nu = r, \text{ at } \partial\Omega.$ <p>3. Calculate $L_{\varepsilon, \alpha, \beta}(\phi_n, \psi_n^1, \psi_n^2)$ and $L_{\varepsilon, \alpha, \beta}^j(\phi_n, \psi_n^1, \psi_n^2)$ as in (24).</p> <p>4. Evaluate the updates $\delta\phi \in H^1(\Omega)$, $\delta\psi^j \in \text{BV}(\Omega)$ by solving (23)</p> <p>5. Update the level set functions $\phi_{n+1} = \phi_n + \frac{1}{\alpha}\delta\phi$, and the level function values $\psi_{n+1}^j = \psi_n^j + \frac{1}{\alpha}\delta\psi^j$.</p> |
|--|

Table 2: An explicit algorithm based on the proposed level set iterative regularization method.

7 Conclusions and Future Directions

In this article, we generalize the results of convergence and stability of the level set regularization approach proposed in [14, 13] to the case where the level values and the level set are non-constant inside of each region. We analyze the particular case, where the set Ω is divide in two regions. It is easy to extend the analysis to the case of multiple regions adapting the multiple level set approach in [15, 14].

We shows the applicability of the level set framework introduced in this paper for an inverse potential problem with piecewise smooth components. The application is a natural generalization of the inverse potential problem computed in [15, 13, 14]. Applications of inverse potential problems in EMG appears in [39, 38]. We also investigate the applicability in the identification of inhomogeneities inside a nonlinear magnetic workpiece. In both application, an iterative algorithm based on the optimality condition of the smooth Tikhonov functional $\mathcal{G}_{\varepsilon, \alpha}$ was proposed.

A natural continuation of this paper is the numerical implementation. Level set numerical implementations for the inverse potential problem was done before in [15, 14, 13], where the level values are assumed to be constant. Implementations of level set methods for resistivity/conductivity problem in elliptic equation have been intensively implemented recently. See, for example, [9, 18, 31, 40, 36, 11, 5] and references.

References

- [1] R. Acar and C. R. Vogel, *Analysis of bounded variation penalty methods for ill-posed problems*, Inverse Problems **10** (1994), no. 6, 1217–1229.
- [2] R.A. Adams, *Sobolev spaces*, Academic Press, New York, 1975.
- [3] P. A. Berthelsen, *A decomposed immersed interface method for variable coefficient elliptic equations with non-smooth and discontinuous solutions*, J. Comput. Phys. **197** (2004), no. 1, 364–386.
- [4] M. Burger, *A level set method for inverse problems*, Inverse Problems **17** (2001), no. 5, 1327–1355.
- [5] M. Burger and S.J. Osher, *A survey on level set methods for inverse problems and optimal design*, European J. Appl. Math. **16** (2005), no. 2, 263–301.
- [6] T.F. Chan and X.-C. Tai, *Identification of discontinuous coefficients in elliptic problems using total variation regularization*, SIAM J. Sci. Comput. **25** (2003), no. 3, 881–904.

- [7] ———, *Level set and total variation regularization for elliptic inverse problems with discontinuous coefficients*, J. Comput. Phys. **193** (2004), no. 1, 40–66.
- [8] F. Chantalat, C-H. Bruneau, C. Galusinski, and A. Iollo, *Level-set, penalization and Cartesian meshes: a paradigm for inverse problems and optimal design*, J. Comput. Phys. **228** (2009), no. 17, 6291–6315.
- [9] E.T. Chung, T.F. Chan, and X.-C. Tai, *Electrical impedance tomography using level set representation and total variational regularization*, J. Comput. Phys. **205** (2005), no. 1, 357–372.
- [10] J. Chung and L. Vese, *Image segmentation using a multilayer level-sets approach*, UCLA C.A.M. Report **193** (2003), no. 03-53, 1–28.
- [11] I. Cimrák and R. Van Keer, *Level set method for the inverse elliptic problem in nonlinear electromagnetism*, J. Comput. Phys. **229** (2010), no. 24, 9269–9283.
- [12] R. Dautray and J.-L. Lions, *Mathematical analysis and numerical methods for science and technology. Vol. 2*, Springer-Verlag, Berlin, 1988.
- [13] A. De Cezaro and A. Leitão, *Level-set of L^2 type for recovering shape and contrast in inverse problems*, Inverse Problems in Science and Engineering (**to appear**) (2011), 1–20.
- [14] A. De Cezaro, A. Leitão, and X.-C. Tai, *On a level-set type methods for recovering piecewise constant solution of ill-posed problems*, Lecture Notes in Computer Science **5567** (2009), 50–62.
- [15] ———, *On multiple level-set regularization methods for inverse problems*, Inverse Problems **25** (2009), no. 035004, 22p.
- [16] M. Delfour and J-P. Zolesio, *Shape analysis via oriented distance functions*, Journal of Functional Analysis **123** (1994), 129–201.
- [17] B. Dong, A. Chien, Y. Mao, J. Ye, F. Vinuela, and S. Osher, *Level set based brain aneurysm capturing in 3D*, Inverse Probl. Imaging **4** (2010), no. 2, 241–255.
- [18] O. Dorn and D. Lesselier, *Level set methods for inverse scattering—some recent developments*, Inverse Problems **25** (2009), no. 12, 125001, 11.
- [19] H. W. Engl, M. Hanke, and A. Neubauer, *Regularization of inverse problems*, Mathematics and its Applications, vol. 375, Kluwer Academic Publishers Group, Dordrecht, 1996.
- [20] H.W. Engl, K. Kunisch, and A. Neubauer, *Convergence rates for Tikhonov regularisation of nonlinear ill-posed problems*, Inverse Problems **5** (1989), no. 4, 523–540.
- [21] L.C. Evans and R.F. Gariepy, *Measure theory and fine properties of functions*, Studies in Advanced Mathematics, CRC Press, Boca Raton, FL, 1992.
- [22] F. Frühauf, O. Scherzer, and A. Leitão, *Analysis of regularization methods for the solution of ill-posed problems involving discontinuous operators*, SIAM J. Numer. Anal. **43** (2005), 767–786.
- [23] F. Hettlich and W. Rundell, *Iterative methods for the reconstruction of an inverse potential problem*, Inverse Problems **12** (1996), no. 3, 251–266.
- [24] V. Isakov, *Inverse source problems*, Mathematical Surveys and Monographs, vol. 34, American Mathematical Society, Providence, RI, 1990.
- [25] A. Leitão and O. Scherzer, *On the relation between constraint regularization, level sets, and shape optimization*, Inverse Problems **19** (2003), L1–L11.

- [26] J. Lie, M. Lysaker, and X.-C. Tai, *A variant of the level set method and applications to image segmentation*, Math. Comp. **75** (2006), no. 255, 1155–1174.
- [27] E. Maitre and F. Santosa, *Level set methods for optimization problems involving geometry and constraints. II. Optimization over a fixed surface*, J. Comput. Phys. **227** (2008), no. 22, 9596–9611.
- [28] N. G. Meyers, *An L^p -estimate for the gradient of solutions of second order elliptic divergence equations*, Ann. Scuola Norm. Sup. Pisa (3) **17** (1963), 189–206.
- [29] B. F. Nielsen, M. Lysaker, and A. Tveito, *On the use of the resting potential and level set methods for identifying ischemic heart disease: an inverse problem*, J. Comput. Phys. **220** (2007), no. 2, 772–790.
- [30] F. Santosa, *A level-set approach for inverse problems involving obstacles*, ESAIM Contrôle Optim. Calc. Var. **1** (1995/96), 17–33.
- [31] M. Soleimani, W.R.B. Lionheart, and O. Dorn, *Level set reconstruction of conductivity and permittivity from boundary electrical measurements using experimental data*, Inverse Problems in Science and Engineering **14** (2006), 193–210.
- [32] X.-C. Tai and T.F. Chan, *A survey on multiple level set methods with applications for identifying piecewise constant functions*, Int. J. Numer. Anal. Model. **1** (2004), no. 1, 25–47.
- [33] X.-C. Tai and H. Li, *A piecewise constant level set method for elliptic inverse problems*, Appl. Numer. Math. **57** (2007), no. 5-7, 686–696.
- [34] Xue-Cheng Tai and Yao Chang-Hui, *Image segmentation by piecewise constant mumford-shah model without estimating the constants*, CAM report **06-18** (2006), 1–9.
- [35] Y.-H. R. Tsai and S. Osher, *Total variation and level set methods in image science*, Acta Numer. **14** (2005), 509–573.
- [36] K. van den Doel and U. M. Ascher, *On level set regularization for highly ill-posed distributed parameter estimation problems*, J. Comput. Phys. **216** (2006), no. 2, 707–723.
- [37] K. van den Doel, U. M. Ascher, and A. Leitão, *Multiple level sets for piecewise constant surface reconstruction in highly ill-posed problems*, J. Sci. Comput. **43** (2010), no. 1, 44–66.
- [38] K. van den Doel, U. M. Ascher, and D. K. Pai, *Computed myography: three-dimensional reconstruction of motor functions from surface EMG data*, Inverse Problems **24** (2008), no. 6, 065010, 17.
- [39] ———, *Source localization in electromyography using the inverse potential problem*, Inverse Problems **27** (2011), no. 2, 025008, 20.
- [40] K. van den Doel and U.M. Ascher, *Dynamic level set regularization for large distributed parameter estimation problems*, Inverse Problems **23** (2007), 1271–1288.
- [41] L.A. Vese and S.J. Osher, *Image denoising and decomposition with total variation minimization and oscillatory functions*, J. Math. Imaging Vision **20** (2004), no. 1-2, 7–18, Special issue on mathematics and image analysis.
- [42] K. Yosida, *Functional analysis*, Classics in Mathematics, Springer-Verlag, Berlin, 1995, Reprint of the sixth (1980) edition.